

Non-commutative Twistor Space

K.C. HANNABUSS

Balliol College, Oxford, OX1 3BJ, England. e-mail: kch@ermine.ox.ac.uk

Mathematics Subject Classifications (2000): 58B34, 83C60, 83C65, 35Q58, 53D55

Keywords: non-commutative geometry, deformation, twistor, integrable model.

Abstract: It is shown that deformations of twistor space compatible with the Moyal deformation of Minkowski space-time must take the form recently suggested by Kapustin, Kuznetsov and Orlov, [4]

0. Introduction

In an important contribution to the burgeoning study of integrable systems in non-commutative geometry (see [3], [6], [8]), Kapustin, Kuznetsov and Orlov have generalised the Penrose-Ward twistor transform to that setting, [4]. The geometric parts of the construction translate fairly naturally using the standard techniques of non-commutative geometry ([2],[5]), but the form (or even existence) of the non-commutative twistor algebra is less obvious. In [4] such an algebra is presented using braiding properties defined by an appropriate R -matrix. In this note we consider the extent to which this is determined by the deformation of the original space-time algebra using a dual quantum group description. The ordinary twistor space and the six-dimensional conformal space in which compactified Minkowski space-time is identified with the rays of a cone are both flag manifolds generated by the conformal group action on highest weight vectors, so we consider non-commutative twistor space as a quantum flag manifold for the quantised function algebra of a deformation of the conformal group [1]. We shall also give a coordinate-free expression for the R -matrix.

1. The non-commutative space-time algebra

We recall that the non-commutative algebra for (four-dimensional) space-time $\mathcal{M}$ is the twisted group algebra defined by a symplectic form θ on $\mathcal{M}$. Its Lie algebra is generated by elements $X(u)$ for $u \in \mathcal{M}$ which satisfy $[X(u), X(v)] = i\hbar\theta(u, v)I$, where I is a positive central element. Using the non-degenerate symmetric bilinear form G on $\mathcal{M}$ we may write $i\hbar\theta(u, v) = G(u, \Theta v)$ for a suitable skew-symmetric operator Θ , (which may be regarded as an element of the Lie algebra of the orthogonal group of G), so that

$$[X(u), X(v)] = G(u, \Theta v)I.$$

We can also define $H = \frac{1}{2}G^{ij}X_iX_j$ (where $X(u) = X_ju^j$ in terms of coordinates and using the summation convention), and calculate that

$$[H, X(u)] = X(\Theta u)I.$$

In order to generalise the conformal theory it is useful to introduce a central element T such that $I = T^2$, and to define $D = T^{-1}H$, so that the commutation relations take the homogeneous form

$$[X(u), X(v)] = G(u, \Theta v)T^2, \quad [D, X(u)] = X(\Theta u)T,$$

defining a quadratic algebra. The identity $2TD = 2H = G^{ij}X_iX_j$, gives the non-commutative version of the quadric cone of the conformal theory.

In the commutative case one studies connections with self-dual curvature. This curvature vanishes on isotropic planes and these define twistors (projective spinors for the conformal group) $\mathcal{T}$, [7]. (For an isotropic plane through the origin, we note that the Clifford algebra of $\mathcal{M}$ acts on the space of spinors. Writing $\gamma(u)$ for the action of $u \in \mathcal{M}$ we may consider the “vacuum” vector $\Psi_\xi \in \mathcal{T}$ with the property that $\gamma(u)\Psi_\xi = 0$ for all u in the plane defined by $\xi \in \wedge^2\mathcal{M}$, that is for $u \wedge \xi = 0$. Since Ψ is determined only up to multiples

this gives a ray, and defines a point of the projective space of $\mathcal{T}$. Isotropic planes through other points can be obtained using the action of the conformal group.)

The space $\mathcal{T}$ decomposes into odd and even spinors $\mathcal{T}_+ \oplus \mathcal{T}_-$. Each of the spaces $\mathcal{T}_\pm$ is two-dimensional so that $\wedge^2 \mathcal{T}_\pm$ is spanned by a single vector $\epsilon_\pm$. (We normalise these so that $\|\epsilon_\pm\|^2 = 2$, so that when we choose orthonormal bases e_1, e_2 for $\mathcal{T}_+$ and e_3, e_4 for $\mathcal{T}_-$ we may take $\epsilon_+ = e_1 \wedge e_2 = e_1 \otimes e_2 - e_2 \otimes e_1$ and $\epsilon_- = e_3 \wedge e_4 = e_3 \otimes e_4 - e_4 \otimes e_3$.)

The six-dimensional space $\wedge^2 \mathcal{T}$ has a natural symmetric bilinear form B defined by

$$B(\xi, \eta) \epsilon_+ \wedge \epsilon_- = \xi \wedge \eta.$$

The action of the conformal group on $\wedge^2 \mathcal{T}$ leaves invariant the null cone $B(u, u) = 0$, and the rays of this cone form a compact complex version of space-time. (The finite points of space time can be identified with the rays of the cone for which $B(\epsilon_-, u) \neq 0$, or equivalently $\mathcal{M}$ can be identified with $\mathcal{T}_+ \wedge \mathcal{T}_-$. For brevity we shall call $\wedge^2 \mathcal{T}$ conformal space-time.)

By construction the orthogonal group of G has its spin representation Γ on $\mathcal{T}$, which leaves each of $\mathcal{T}_\pm$ invariant, and so the forms $\epsilon_\pm$ are also invariant under the spin action. We may choose the orthonormal bases e_1, e_2 for $\mathcal{T}_+$ and e_3, e_4 for $\mathcal{T}_-$ so that the matrix $\Gamma(\Theta)$ is diagonal:

$$\frac{1}{2} \hbar \begin{pmatrix} (b-a)\sigma_3 & 0 \\ 0 & (a+b)\sigma_3 \end{pmatrix} \quad \text{where} \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

We now complete $\epsilon_\pm$ to a basis for $\wedge^2 \mathcal{T}$ by introducing $\epsilon_1 + i\epsilon_4 = e_4 \wedge e_1$, $\epsilon_1 - i\epsilon_4 = e_2 \wedge e_3$, $\epsilon_2 + i\epsilon_3 = e_1 \wedge e_3$, and $\epsilon_2 - i\epsilon_3 = e_2 \wedge e_4$. With this notation $\{\epsilon_\mu\}$ forms a G -orthonormal basis of $\mathcal{M} \sim \wedge^2 \mathcal{T}$ when we identify G with $-2B$. The matrix of Θ with respect to this basis is given by

$$i\hbar \begin{pmatrix} 0 & 0 & 0 & a \\ 0 & 0 & b & 0 \\ 0 & -b & 0 & 0 \\ -a & 0 & 0 & 0 \end{pmatrix}.$$

Unfortunately, the commutation relations of the non-commutative theory destroy the full conformal symmetry. For example, the transformations of the space $\mathcal{M}$, affecting only the operators $X(u)$, must not only be orthogonal with respect to G but also be symplectic with respect to θ , or equivalently commute with Θ . In general this forces them to lie in a Cartan subgroup, though, for particular θ , (when a vanishes) they may lie in a larger group. We must therefore deform the conformal group, or more precisely we shall quantise its function algebra by deforming the R matrix associated with the commutation relations. To make explicit the connection between commutation relations and an R matrix, we consider for any vector space U an operator $R : U \otimes U \rightarrow U \otimes U$ satisfying the Yang-Baxter equation, and write $R = \Phi + R'$ where Φ denotes the usual flip operator $x \otimes y \mapsto y \otimes x$ on $U \otimes U$, and R' is the deformation. The function algebra defined by R is given by operators A on U which satisfy $R(A \otimes A) = (A \otimes A)R$. The following is then a straightforward consequence of our definitions.

PROPOSITION 1. The relations $R(A \otimes A) = (A \otimes A)R$ of the function algebra defined by $R = \Phi + R'$ are equivalent to the commutation relations

$$[\langle e|Au\rangle, \langle f|Av\rangle] = \langle f \otimes e|(A^{(2)}R' - R'A^{(2)})(u \otimes v)\rangle,$$

where $A^{(2)} = A \otimes A$, $e, f \in U^*$, $u, v \in U$, and $\langle e|u\rangle$ denotes the pairing between the dual spaces U^* and U .

Proof: The relations $R(A \otimes A) = (A \otimes A)R$ can be rewritten as

$$\Phi(A \otimes A) - (A \otimes A)\Phi = (A \otimes A)R' - R'(A \otimes A).$$

Writing $A^{(2)} = A \otimes A$, and choosing $e, f \in U^*$, $u, v \in U$ this gives

$$\langle f \otimes e|(\Phi A^{(2)} - A^{(2)}\Phi)(u \otimes v)\rangle = \langle f \otimes e|(A^{(2)}R' - R'A^{(2)})(u \otimes v)\rangle.$$

The left-hand side can be written as

$$\langle e \otimes f | A^{(2)} u \otimes v \rangle - \langle f \otimes e | A^{(2)} v \otimes u \rangle = \langle e | Au \rangle \langle f | Av \rangle - \langle f | Av \rangle \langle e | Au \rangle = [\langle e | Au \rangle, \langle f | Av \rangle],$$

so that we have the stated result. $\blacksquare$

Let us take $U = \wedge^2 \mathcal{T}$ and write the vectors as $v = v_+ \epsilon_+ + v_0 + v_- \epsilon_-$, with $v_0 \in \mathcal{T}_+ \wedge \mathcal{T}_- \sim \mathcal{M}$. To avoid confusion with the twistor examples we write $\tilde{R}$ and $\tilde{A}$ instead of R and A . We are less interested in the whole function algebra than in the quantum flag manifold generated by the highest weight vector ϵ_+^* of the dual basis of U^* , and want to identify $\langle \epsilon_+^* | \tilde{A} v \rangle$ with $v_+ D + X(v_0) + v_- T$. In this notation the commutation relations reduce to

$$[u_+ D + X(u_0) + u_- T, v_+ D + X(v_0) + v_- T] = u_+ T X(\Theta v_0) - v_+ X(\Theta u_0) T + G(u_0, \Theta v_0) T^2.$$

One simple way of obtaining these is to consider the deformation given by $\tilde{R}'$

$$\tilde{R}'(u \otimes v) = u_+ \epsilon_- \otimes \Theta v_0 - \Theta u_0 \otimes v_+ \epsilon_- + G(u_0, \Theta v_0) \epsilon_- \otimes \epsilon_-.$$

Since this has no terms in ϵ_+ , we have $\langle \epsilon_+^* \otimes \epsilon_+^* | \tilde{R}' = 0$ and one may readily check that $\tilde{R}$ gives the correct commutation relation. It is useful to note that

$$\tilde{R}'(u \wedge v) = u_+ \epsilon_- \otimes_S \Theta v_0 - \Theta u_0 \otimes_S v_+ \epsilon_- + G(u_0, \Theta v_0) \epsilon_- \otimes_S \epsilon_-,$$

and

$$\tilde{R}'(u \otimes_S v) = u_+ \epsilon_- \wedge \Theta v_0 - \Theta u_0 \wedge v_+ \epsilon_-,$$

so that $\tilde{R}'$ interchanges symmetric and antisymmetric tensors. It is also easy to check that the square of $\tilde{R}'$ vanishes. These are in accord with the following simple result.

PROPOSITION 2. A family of deformations $R = \Phi + R'$ satisfies $R^2 = 1$, if and only if R' maps the symmetric tensors to antisymmetric tensors and vice versa, and $R'^2 = 0$.

Proof: This is a matter of comparing terms of first and second order in R' in the identity

$$1 = R^2 = \Phi^2 + \Phi R' + R' \Phi + R'^2 = 1 + \Phi R' + R' \Phi + R'^2,$$

which gives $\Phi R' + R' \Phi = 0 = R'^2$. The first identity can be rewritten in the form

$$R'(1 \pm \Phi) = (1 \mp \Phi)R',$$

which, recalling that $\frac{1}{2}(1 + \Phi)$ is the projection onto the symmetric tensor product $\otimes_S^2 U$ and $\frac{1}{2}(1 - \Phi)$ projects onto the antisymmetric tensor product $\wedge^2 U$, shows that R' maps $\otimes_S^2 U$ to $\wedge^2 U$ and $\wedge^2 U$ to $\otimes_S^2 U$. (For similar reasons R' also interchanges $\otimes_{S,R}^2 U = \frac{1}{2}(1 + R)(U \otimes U)$ and $\wedge_R^2 U = \frac{1}{2}(1 - R)(U \otimes U)$.) $\blacksquare$

2. Yang-Baxter operators for the twistors

We now want to find a Yang-Baxter operator R for twistor space which is consistent with that on conformal space-time $\wedge^2 \mathcal{T}$. More precisely, a Yang-Baxter operator R on $\otimes^2 \mathcal{T}^*$ can be extended to a Yang-Baxter operator $\tilde{R} : \otimes^4 \mathcal{T} \rightarrow \otimes^4 \mathcal{T}$, defined by

$$\tilde{R} = R_{23} R_{12} R_{34} R_{23}.$$

This interchanges the first and second pairs of factors in the tensor product, allowing one to extend the Yang-Baxter operator R for twistor space to a Yang-Baxter operator $\tilde{R}$ for the conformal space-time $\wedge_R^2 \mathcal{T}$,

and, for consistency, this should agree with that found above. The primary task is to see whether any such R exist, and to find them.

It is easy to see that $\tilde{R}$ inherits various properties of R , but in the Appendix we show that two crucial properties of $\tilde{R}$ must be shared by R , a fact which substantially simplifies the task of finding it.

THEOREM 3. If a family of Yang-Baxter operators R which is a deformation of the flip Φ satisfies $[R_{23}R_{12}R_{34}R_{23}]^2 = 1$, then $R^2 = 1$, and if, in addition, $R_{23}R_{12}R_{34}R_{23}$ commutes with $\Lambda^{(4)} = \Lambda \otimes \Lambda \otimes \Lambda \otimes \Lambda$ for invertible Λ , then R commutes with $\Lambda^{(2)} = \Lambda \otimes \Lambda$.

We have already observed that for conformal space-time $\tilde{R}^2 = 1$, and the invariance of the commutation relations under the infinitesimal rotation Θ means that $\tilde{R}$ commutes with Θ and so with the invertible operator $\Lambda = \exp(s\Theta)$. This suggests that the non-commutative twistor algebra is defined by an operator R satisfying $R^2 = 1$, and commuting with the spin action of $\Lambda \otimes \Lambda = \exp(s\Theta_1 + s\Theta_2)$, where $\Theta_1 = \Gamma(\Theta) \otimes 1$, and $\Theta_2 = 1 \otimes \Gamma(\Theta)$. (It is merely suggested because the space-time $\tilde{R}$ is only defined on the subspace $\wedge_R^2 \mathcal{T}^* \otimes \wedge_R^2 \mathcal{T}^*$. There is no problem provided that $\tilde{R}$ has an extension to the whole of $\otimes^4 \mathcal{T}^*$ with the same involutory and intertwining properties. Our argument will start by assuming that there is such an extension, and then justifying this once R has been found. Once established we can deduce that R also commutes with the spin action $\Theta_1 + \Theta_2$.) In fact $\tilde{R}$ enjoys an additional, less obvious symmetry arising from its compatibility with the exact sequence: $0 \longrightarrow \mathcal{T}_- \longrightarrow \mathcal{T} \longrightarrow \mathcal{T}_+ \longrightarrow 0$.

PROPOSITION 4. Let N be a nilpotent operator on $\mathcal{T}$ such that $\text{im}(N) = \ker(N) = \mathcal{T}_-$ and let $K_s = 1 + sN$. Then $K_s^{(4)} = K_s \otimes K_s \otimes K_s \otimes K_s$ commutes with $\tilde{R}$.

Proof: By definition K_s acts as the identity on $\mathcal{T}_-$ and so $K_s^{(2)}\epsilon_- = \epsilon_-$. By contrast $K_s^{(2)}\epsilon_+$ must have the general form $\epsilon_+ + sn + s^2\nu\epsilon_-$ for scalar ν and $n \in \mathcal{T}_+ \wedge \mathcal{T}_-$. For $u_0 \in \mathcal{T}_+ \wedge \mathcal{T}_-$ we must have $K_s^{(2)}u_0 = u_0 + s\lambda\epsilon_-$ for a suitable scalar λ which we shall now find. We first notice that the determinant of K_s is clearly 1, so that for any ξ and $\eta \in \mathcal{T} \wedge \mathcal{T}$ we have $\xi \wedge \eta = K_s^{(2)}\xi \wedge K_s^{(2)}\eta$, and thence $B(\xi, \eta) = B(K_s^{(2)}\xi, K_s^{(2)}\eta)$. In particular, taking $\xi = \epsilon_+$ and $\eta = u_0$ gives

$$0 = B(\epsilon_+, u_0) = B(\epsilon_+ + sn + s^2\nu\epsilon_-, u_0 + s\lambda\epsilon_-) = s\lambda + sB(n, u_0),$$

so that $\lambda = -B(n, u_0) = \frac{1}{2}G(n, u_0)$, and $K_s^{(2)}u_0 = u_0 + \frac{1}{2}sG(n, u_0)\epsilon_-$.

We now calculate that

$$\begin{aligned} K_s^{(4)}\tilde{R}'(u \otimes v) &= K_s^{(4)}(u_+\epsilon_- \otimes \Theta v_0 - \Theta u_0 \otimes v_+\epsilon_- + G(u_0, \Theta v_0)\epsilon_- \otimes \epsilon_-) \\ &= u_+\epsilon_- \otimes (\Theta v_0 + \frac{1}{2}sG(n, \Theta v_0)\epsilon_- - (\Theta u_0 + \frac{1}{2}sG(n, \Theta u_0)\epsilon_-) \otimes v_+\epsilon_- + G(u_0, \Theta v_0)\epsilon_- \otimes \epsilon_-). \end{aligned}$$

Since $G(n, \Theta u_0) = -G(u_0, \Theta n)$ and, in particular, $G(n, \Theta n) = 0$, this reduces to

$$u_+\epsilon_- \otimes \Theta v_0 - \Theta u_0 \otimes v_+\epsilon_- + G(u_0 + \frac{1}{2}su_+n, \Theta(v_0 + \frac{1}{2}sv_+n))\epsilon_- \otimes \epsilon_- = \tilde{R}'(K_s^{(4)}(u \otimes v)),$$

and since $\tilde{\Phi}$ automatically commutes with any operator $\Lambda^{(4)}$ this gives the stated result. $\blacksquare$

We also want R to satisfy the Yang-Baxter equation, but there is another requirement which is much stronger, namely that the restriction of $\tilde{R} = R_{23}R_{12}R_{34}R_{23}$ to $\wedge_R^2 \mathcal{T}^* \otimes \wedge_R^2 \mathcal{T}^*$ contains only first order terms in Θ (that is, in R'). With the requirements that R be an involution and have the same symmetries as $\tilde{R}$, this limits the possibilities considerably.

THEOREM 5. For non-degenerate Θ , any Yang-Baxter operator $R = \Phi + R'$ satisfying $R^2 = 1$, commuting with Θ and giving only first order corrections to $\tilde{R}$ on the conformal spacetime is in the two-parameter family

$$R' = \alpha\Theta_1|\epsilon_- \rangle \langle \epsilon_+^*| + \beta|\epsilon_- \rangle \langle \epsilon_+^*|\Theta_1,$$

for some scalars $\alpha, \beta \in \mathbb{C}$. These operators automatically satisfy the Yang-Baxter equation.

Proof: Since R' commutes with the action of Θ it must preserve eigenspaces for the action. The formula for $\Gamma(\Theta)$ shows that generically (unless a , b or $a \pm b$ vanishes) each symmetric tensor $e_j \otimes e_j$ is in a one-dimensional eigenspace of its own, and since R' maps symmetric tensors to antisymmetric tensors we must have $e_j \otimes e_j \in \ker(R')$. However, $\ker(R')$ must be invariant for operators of the form $K_s^{(2)}$ and by suitable choices of N , $K_s e_j \otimes K_s e_j$ ($j = 1, 2$) generate the whole of $\mathcal{T} \otimes_S \mathcal{T}_-$ which must therefore also lie in the kernel.

Generically, there is also a four-dimensional null space for the action of Θ spanned by $e_1 \otimes_S e_2$, $\epsilon_+ = e_1 \wedge e_2$, $e_3 \otimes_S e_4$, and $\epsilon_- = e_3 \wedge e_4$, and the other eigenspaces are two-dimensional spanned by tensors of the form $e_j \otimes_S e_k$ and $e_j \wedge e_k$ for various distinct $\{j, k\} \notin \{\{1, 2\}, \{3, 4\}\}$. We have just seen that the symmetric tensors are in the kernel of R' . Since R' maps antisymmetric to symmetric tensors each $e_j \wedge e_k$ must map to a multiple of $e_j \otimes_S e_k$. In fact, this multiple must also vanish otherwise overlapping terms such as $R'_{12} R'_{23}(e_i \otimes e_j \otimes e_k)$, give quadratic contributions to $\tilde{R}$, so we may assume that $R'(e_j^* \otimes_S e_k^*) = 0$ and $R'(e_j^* \wedge e_k^*) = 0$ for the given range of j and k .

The only interesting deformations therefore occur in the four-dimensional null space. Here it is useful to note that $\Theta_1 \epsilon_+ = \frac{1}{2} \hbar(b - a) e_1 \otimes_S e_2$, and $\Theta_1 \epsilon_- = \frac{1}{2} \hbar(b + a) e_3 \otimes_S e_4$. (Since $\epsilon_{\pm}$ and their duals are invariant under the appropriate actions of $\Theta_1 + \Theta_2$, these could equally have been written in terms of Θ_2 .) We already know that R' kills $e_3 \otimes_S e_4$ and $e_3 \wedge e_4$, so, again discarding the deformations which mix the symmetric and antisymmetric products of the same vectors and lead to quadratic contributions to $\tilde{R}$, we are left with R' of the form

$$R' \epsilon_+ = \alpha \Theta_1 \epsilon_- \quad R' \Theta_1 \epsilon_+ = \alpha' \epsilon_-,$$

or, equivalently, $R' = \alpha \Theta_1 | \epsilon_- \rangle \langle \epsilon_+^* | + \beta | \epsilon_- \rangle \langle \epsilon_+^* | \Theta_1$. The condition that $R'^2 = 0$ holds automatically as do the Yang-Baxter equations. ■

We shall next show that with an appropriate choice of constants α and β this is compatible with $\tilde{R}$. Since by construction $R_{23} R_{12} R_{34} R_{23}$ is an involution commuting with the same operators as $\tilde{R}$ this verifies that there is indeed an extension of $\tilde{R}$ to $\otimes^4 \mathcal{T}^*$, so justifying our working assumption.

THEOREM 6. The only Yang-Baxter operator for $\mathcal{T}$ compatible with the known Yang-Baxter operator $\tilde{R}$ on the non-commutative conformal space-time is

$$R = \Phi + 2[\Theta_1, | \epsilon_+^* \rangle \langle \epsilon_- |]_+.$$

Proof: We have noted that there are only first order corrections so the Yang-Baxter operator takes the form $\tilde{R} = \tilde{\Phi} + \tilde{R}'$, where $\tilde{\Phi}$ is the flip and

$$\tilde{R}' = \Phi_{23} \Phi_{12} \Phi_{34} R'_{23} + \Phi_{23} \Phi_{12} R'_{34} \Phi_{23} + \Phi_{23} R'_{12} \Phi_{34} \Phi_{23} + R'_{23} \Phi_{12} \Phi_{34} \Phi_{23}.$$

Since, for example $\Phi_{23} R_{24} = R'_{34} \Phi_{23}$, this can be rewritten as

$$\tilde{R}' = R'_{14} \Phi_{23} \Phi_{12} \Phi_{34} + R'_{24} \Phi_{13} + R'_{13} \Phi_{24} + R'_{23} \Phi_{12} \Phi_{34} \Phi_{23}.$$

Applying this to $(u \wedge x) \otimes (v \wedge y)$ we get

$$\begin{aligned} & R'_{14}(x \otimes y \otimes u \otimes v - u \otimes y \otimes x \otimes v - x \otimes v \otimes u \otimes y + u \otimes v \otimes x \otimes y) \\ & + R'_{24}(v \otimes x \otimes u \otimes y - v \otimes u \otimes x \otimes y - y \otimes x \otimes u \otimes v + y \otimes u \otimes x \otimes v) \\ & + R'_{13}(u \otimes y \otimes v \otimes x - x \otimes y \otimes u \otimes v - u \otimes v \otimes y \otimes x + x \otimes v \otimes y \otimes u) \\ & + R'_{23}(v \otimes u \otimes y \otimes x - v \otimes x \otimes y \otimes u - y \otimes u \otimes v \otimes x + y \otimes x \otimes u \otimes v). \end{aligned}$$

If we take $u, v \in \mathcal{T}_+$ and $x, y \in \mathcal{T}_-$ and use the form of R' most of these terms disappear, leaving

$$-R'_{14} u \otimes y \otimes x \otimes v + R'_{24} y \otimes u \otimes x \otimes v + R'_{13} u \otimes y \otimes v \otimes x - R'_{23} y \otimes u \otimes v \otimes x.$$

It will be sufficient to consider the case of $u = e_1$ and $v = e_2$, where

$$\begin{aligned} R(e_1 \otimes e_2) &= \frac{1}{2}\hbar(a+b)\alpha e_3 \otimes_S e_4 + \frac{1}{2}\hbar(b-a)\beta e_3 \wedge e_4 \\ &= \frac{1}{2}\hbar(\alpha(a+b) + \beta(b-a))e_3 \otimes e_4 + \frac{1}{2}\hbar(\alpha(a+b) - \beta(b-a))e_4 \otimes e_3. \end{aligned}$$

Writing $\sigma = \frac{1}{2}\hbar(\alpha(a+b) + \beta(b-a))$ and $\delta = \frac{1}{2}\hbar(\alpha(a+b) - \beta(b-a))$, we reduce $\tilde{R}'(e_1 \wedge x \otimes e_2 \wedge y)$ to

$$\begin{aligned} &\sigma(-e_3 \otimes y \otimes x \otimes e_4 + y \otimes e_3 \otimes x \otimes e_4 + e_3 \otimes y \otimes e_4 \otimes x - y \otimes e_3 \otimes e_4 \otimes x) \\ &+ \delta(-e_4 \otimes y \otimes x \otimes e_3 + y \otimes e_4 \otimes x \otimes e_3 + e_4 \otimes y \otimes e_3 \otimes x - y \otimes e_4 \otimes e_3 \otimes x) \\ &= \sigma y \wedge e_3 \otimes x \wedge e_4 + \delta y \wedge e_4 \otimes x \wedge e_3. \end{aligned}$$

This is non-vanishing if $x = e_3$ and $y = e_4$, or vice versa. The first possibility gives

$$\tilde{R}'(e_1 \wedge x \otimes e_2 \wedge y) = \sigma e_4 \wedge e_3 \otimes e_3 \wedge e_4 = -\sigma \epsilon_- \otimes \epsilon_-,$$

whilst the second gives

$$\tilde{R}'(e_1 \wedge e_4 \otimes e_2 \wedge e_3) = -\delta \epsilon_- \otimes \epsilon_-.$$

On the other hand $e_1 \wedge e_3 = \epsilon_2 + i\epsilon_3$ and $e_2 \wedge e_4 = \epsilon_2 - i\epsilon_3$ gives

$$\tilde{R}'(e_1 \wedge e_3 \otimes e_2 \wedge e_4) = G(e_1 \wedge e_3, \Theta(e_2 \wedge e_4))\epsilon_- \otimes \epsilon_- = -2\hbar b \epsilon_- \otimes \epsilon_-.$$

This gives $\frac{1}{2}(\alpha(a+b) + \beta(b-a)) = 2b$, so that we take $\alpha = \beta = 2$. This is also the choice appropriate to the other choice of x and y and leads to the stated solution, which expressed in terms of coordinates agrees with that of [4]. We can also check that this form of R' works on tensors of the form $\epsilon_+ \otimes u_0$. However, at this point we must mention a subtlety which has so far been suppressed, that we should really be working not with $\wedge^2 \mathcal{T}$ but with $\wedge_R^2 \mathcal{T} = (1 - R) \otimes^2 (\mathcal{T})$. Fortunately, the correction R' has a large kernel, and the only difference is for multiples of ϵ_+ , which should be replaced by $\frac{1}{2}(1 - R)\epsilon_+ = \epsilon_+ - \Theta_1 \epsilon_-$. Fortunately this does not cause any serious complications. ■

3. Commutation relations for twistors

We can immediately combine Theorem 6 with Proposition 1 to give an explicit form for the twistor commutation relations.

COROLLARY 7. The commutation relations for non-commutative twistor space can be written as

$$\begin{aligned} \frac{1}{2}[\langle e|Au\rangle, \langle f|Av\rangle] &= \langle f \otimes e|A^{(2)}\Theta_1\epsilon_-\rangle\langle \epsilon_+^*|u \otimes v\rangle + \langle f \otimes e|A^{(2)}\epsilon_-\rangle\langle \epsilon_+^*|\Theta_1(u \otimes v)\rangle \\ &\quad - \langle f \otimes e|\Theta_1\epsilon_-\rangle\langle \epsilon_+^*|Au \otimes Av\rangle - \langle f \otimes e|\epsilon_-\rangle\langle \epsilon_+^*|\Theta_1(Au \otimes Av)\rangle. \end{aligned}$$

Recalling that $\ker(R') \supseteq \mathcal{T}_- \otimes \mathcal{T} + \mathcal{T} \otimes \mathcal{T}_-$, we see that the first term on the right-hand side vanishes whenever u or v lies in $\mathcal{T}_-$. Similarly, the fact that $\text{im}(R') \subseteq \mathcal{T}_- \otimes \mathcal{T}_-$ means that the second term on the right vanishes for e or f in $\mathcal{T}_+^*$. Taken together we see that $[\langle e|Au\rangle, \langle f|Av\rangle] = 0$ whenever e or $f \in \mathcal{T}_+^*$ and u or $v \in \mathcal{T}_-$, and, in particular, elements of the form $\langle e|Au\rangle$ with $e \in \mathcal{T}_+^*$ and $u \in \mathcal{T}_-$ are central in the algebra.

More generally it is sufficient for our purposes to use only e and f in $\mathcal{T}_+^*$, and then we derive the commutation relation

$$\frac{1}{2}[\langle e|Au\rangle, \langle f|Av\rangle] = \langle f \otimes e|A^{(2)}\Theta_1\epsilon_-\rangle\langle \epsilon_+^*|u \otimes v\rangle + \langle f \otimes e|A^{(2)}\epsilon_-\rangle\langle \epsilon_+^*|\Theta_1(u \otimes v)\rangle.$$

Introducing the symplectic forms $\omega_\pm(u, v) = \langle \epsilon_\pm|u \otimes v\rangle$ on $\mathcal{T}_\pm$, and using the self-adjointness of Θ_1 , we may rewrite this as

$$\begin{aligned} \frac{1}{2}[\langle e|Au\rangle, \langle f|Av\rangle] &= \langle f \otimes e|A^{(2)}\Theta_1\epsilon_-\rangle\omega_+(u, v) + \langle f \otimes e|A^{(2)}\epsilon_-\rangle\omega_+(\Gamma(\Theta)u, v) \\ &= \langle f \otimes e|A^{(2)}[\omega_+(u, v)\Theta_1 + \omega_+(\Gamma(\Theta)u, v)]\epsilon_-\rangle. \end{aligned}$$

The most interesting case arises for $u = e_1$ and $v = e_2$ when $\omega_+(u, v) = 1$ and $\omega_+(\Gamma(\Theta)u, v) = \frac{1}{2}\hbar(b - a)$, so that

$$\begin{aligned} [\langle e|Ae_1\rangle, \langle f|Ae_2\rangle] &= \hbar\langle f \otimes e|A^{(2)}((b+a)e_3 \otimes_S e_4 + (b-a)e_3 \wedge e_4)\rangle \\ &= 2\hbar\langle f \otimes e|A^{(2)}(be_3 \otimes e_4 + ae_4 \otimes e_3)\rangle. \end{aligned}$$

Setting $A_{jr} = \langle e_j|Ae_r\rangle$ and taking $e = e_r$ and $f = e_s$, we deduce that whenever r and s are 1 or 2, the commutation relation gives

$$[A_{r1}, A_{s2}] = 2\hbar(bA_{s3}A_{r4} + aA_{s4}A_{r3}).$$

Setting $z_j = A_{1j}$, $w_j = A_{2j}$ and recalling that we already know that z_3, z_4, w_3 , and w_4 are central, we obtain the relations

$$[z_1, z_2] = 2\hbar(a+b)z_3z_4 \quad [z_1, w_2] = 2\hbar(az_3w_4 + bz_4w_3) \quad [w_1, w_2] = 2\hbar(a+b)w_3w_4.$$

The first of these is essentially the relation of [4] for the twistor algebra. These provide an alternative way of constructing the Minkowski algebra by rewriting the matrix elements $\langle \epsilon_+^*|A^{(2)}\xi\rangle$ for $\xi \in \wedge_R^2 \mathcal{T}^*$ in terms of $z_{jk} = w_jz_k - w_kz_j$, we get the formulae of [4]. For example, $T = \langle \epsilon_+^*|A^{(2)}\epsilon_-\rangle = z_3w_4 - w_3z_4$ is central to the algebra.

Appendix

This Appendix is devoted to a proof of Theorem 3. This is done in stages through a chain of propositions. We start by rewriting the conditions that $\tilde{R}^2 = 1$, and that $\tilde{R}$ commutes with $\Lambda^{(4)} = \Lambda \otimes \Lambda \otimes \Lambda \otimes \Lambda$. The first condition is equivalent to $\tilde{R} = \tilde{R}^{-1} = [R^{-1}]^\sim$, and the second to $\tilde{R} = \Lambda^{(4)}\tilde{R}\Lambda^{(4)-1} = [\Lambda^{(2)}R\Lambda^{(2)-1}]^\sim$. We should like to see whether these two statements, together with the Yang-Baxter equations, are sufficient to give $R^{-1} = R = \Lambda^{(2)}R\Lambda^{(2)-1}$.

To this end we ask whether two Yang-Baxter operators P and Q satisfying

$$P_{23}P_{12}P_{34}P_{23} = Q_{23}Q_{12}Q_{34}Q_{23}$$

must necessarily be identical, that is $P = Q$.

LEMMA 8. Yang-Baxter operators P and Q satisfy $P_{23}P_{12}P_{34}P_{23} = Q_{23}Q_{12}Q_{34}Q_{23}$ if and only if there exist Y and Z such that

$$\begin{aligned} P_{23}P_{12} &= Q_{23}Q_{12}Z_{23}, & Q_{23}Q_{12} &= Z_{12}P_{23}P_{12}, \\ P_{12}P_{23} &= Q_{12}Q_{23}Y_{12}, & Q_{12}Q_{23} &= Y_{23}P_{12}P_{23}. \end{aligned}$$

Proof: Our main tool is separation of variables, which we shall use repeatedly. We start with our assumed identity, which can be rearranged as

$$Q_{12}^{-1}Q_{23}^{-1}P_{23}P_{12} = Q_{34}Q_{23}P_{23}^{-1}P_{34}^{-1}.$$

Since the left hand side is independent of 4 and the right is independent of 1, we see that each side must act as the identity on those factors and so can be written in the form Z_{23} . Rearranging this gives

$$P_{23}P_{12} = Q_{23}Q_{12}Z_{23}, \quad Q_{34}Q_{23} = Z_{23}P_{34}P_{23},$$

and by dropping the indices by 1 the second equation becomes

$$Q_{23}Q_{12} = Z_{12}P_{23}P_{12}.$$

Similarly, since P_{12} and P_{34} commute (and similarly for Q) the assumed identity can be rearranged as $Q_{34}^{-1}Q_{23}^{-1}P_{23}P_{34} = Q_{12}Q_{23}P_{23}^{-1}P_{12}^{-1}$, and setting both sides equal to Y_{23} gives (with an index shift)

$$P_{12}P_{23} = Q_{12}Q_{23}Y_{12}, \quad Q_{12}Q_{23} = Y_{23}P_{12}P_{23}.$$

It is clearly possible to recover the original equation by eliminating either Y or Z . $\blacksquare$

LEMMA 9. Yang-Baxter operators P and Q satisfy $P_{23}P_{12}P_{34}P_{23} = Q_{23}Q_{12}Q_{34}Q_{23}$ if and only if there exist Y , Z , C and D satisfying

$$Q_{12}Z_{12} = C_2P_{12}, \quad Q_{12}Y_{12} = C_1P_{12}.$$

$$Z_{12}P_{12} = Q_{12}D_1, \quad Y_{12}P_{12} = Q_{12}D_2,$$

and $P_{12}P_{23}P_{12} = Q_{12}Q_{23}Q_{12}D_2 = C_2^{-1}Q_{12}Q_{23}Q_{12}$.

Proof: Using the Yang-Baxter equation for Q and the second equation of each pair in Lemma 8 we obtain

$$Q_{12}Z_{12}P_{23}P_{12} = Q_{12}Q_{23}Q_{12} = Q_{23}Q_{12}Q_{23} = Q_{23}Y_{23}P_{12}P_{23}.$$

Now using the Yang-Baxter equation for P , we deduce that

$$Q_{12}Z_{12}P_{12}^{-1} = Q_{23}Y_{23}P_{23}^{-1},$$

and, since one side is the identity on the first factor and the other on the third, we deduce that both sides are of the form C_2 for some operator C . Rearranging this gives

$$Q_{12}Z_{12} = C_2P_{12}, \quad Q_{23}Y_{23} = C_2P_{23},$$

and by lowering the indices the second equation gives

$$Q_{12}Y_{12} = C_1P_{12}.$$

Similarly using the first equation of each pair

$$Q_{23}Q_{12}Z_{23}P_{23} = P_{23}P_{12}P_{23} = P_{12}P_{23}P_{12} = Q_{12}Q_{23}Y_{12}P_{12},$$

from which we see that

$$Q_{23}^{-1}Z_{23}P_{23} = Q_{12}^{-1}Y_{12}P_{12},$$

and both sides must be of the form D_2 , giving

$$Z_{12}P_{12} = Q_{12}D_1, \quad Y_{12}P_{12} = Q_{12}D_2.$$

Substituting these expressions into the previous identities we obtain $P_{12}P_{23}P_{12} = Q_{12}Q_{23}Q_{12}D_2$, and similarly $P_{12}P_{23}P_{12} = C_2^{-1}Q_{12}Q_{23}Q_{12}$. The argument easily reverses. $\blacksquare$

The next two results cover the cases of real interest to us. For these we limit ourselves to Yang-Baxter operators P and Q which are deformations of the standard flip operators, from which one readily checks that Y , Z , C and D are all deformations of the identity operator. The main observation which we need is that each deformation of the identity operator, such as C , has a unique square root which is also a deformation of the identity. It is given by the binomial expansion of $\sqrt{1+x}$ with $C-1$ substituted for x . (In fact, since $\mathcal{T}$ is finite-dimensional, this can be rewritten as a polynomial in C).

LEMMA 10. If Yang-Baxter operators P and Q satisfy $P_{23}P_{12}P_{34}P_{23} = Q_{23}Q_{12}Q_{34}Q_{23}$ and $Q = P^{-1}$, then $Q = P$, that is $P^2 = 1$.

Proof: When $Q = P^{-1}$ we immediately get the equations

$$(P_{12}P_{23}P_{12})^2 = D_2 = C_2^{-1},$$

so that C and D are inverses. Now we also have

$$D_2 = (P_{12}P_{23}P_{12})^2 = (P_{12}P_{23}P_{12})(P_{23}P_{12}P_{23}) = (P_{12}P_{23})^3,$$

and similarly

$$D_2 = (P_{23}P_{12})^3.$$

Consequently

$$P_{12}D_2 = P_{12}(P_{23}P_{12})^3 = (P_{12}P_{23})^3P_{12} = D_2P_{12},$$

so that D_2 , and therefore C_2 , commute with P_{12} . They similarly commute with P_{23} , which, by lowering the indices, means that D_1 and C_1 commute with P_{12} . (Since C_j clearly commutes with P_{12} for $j > 2$ this shows that all the C_j and P_{rs} commute.)

The remaining equations of Lemma 9 now give

$$Q_{12}^2D_1 = Q_{12}Z_{12}P_{12} = C_2P_{12}^2,$$

and

$$Q_{12}^2D_2 = Q_{12}Y_{12}P_{12} = C_1P_{12}^2.$$

Given that C and D are inverse and commute with P_{12} , these reduce to $P_{12}^4 = C_1C_2$, or $P^4 = C \otimes C$. As noted above C has a unique square root K deformed from the identity, and given that the undeformed value of P^2 is 1 we take square roots to see that $P^2 = K \otimes K$.

Since K is given by a polynomial in C it enjoys the same commutation properties as C , and in particular K_j commutes with each of the P_{rs} . This means that

$$\begin{aligned} K_2^{-2} &= C_2^{-1} \\ &= (P_{12}P_{23}P_{12})^2 \\ &= P_{12}P_{23}P_{12}^2P_{23}P_{12} \\ &= P_{12}P_{23}K_1K_2P_{23}P_{12} \\ &= K_1K_2P_{12}P_{23}^2P_{12} \\ &= K_1K_2^2K_3P_{12}^2 \\ &= K_1^2K_2^3K_3. \end{aligned}$$

Separating the indices we see that $K = 1$, and so $P^2 = K \otimes K = 1$. ■

We now look at the second case where $Q_{12} = \Lambda^{(2)}P_{12}\Lambda^{(2)-1}$. and assume that $P^2 = 1 = Q^2$.

LEMMA 11. If Yang-Baxter operators P and Q satisfy $P_{23}P_{12}P_{34}P_{23} = Q_{23}Q_{12}Q_{34}Q_{23}$ and $P^2 = 1 = Q^2$ then $P = Q$.

Proof: Since P and Q are involutions our earlier identity $C_2P_{12}^2 = Q_{12}^2D_1$ reduces to $C_2 = D_1$, so that C and D must both be multiples of 1, indeed the same multiple, $\lambda 1$. The equation

$$Q_{12}Q_{23}Q_{12}D_2 = C_2^{-1}Q_{12}Q_{23}Q_{12}.$$

then shows that $\lambda^2 = 1$, and using the undeformed value of $C = 1$ we deduce that $C = 1 = D$. This means that

$$\begin{aligned} Q_{12}Z_{12} &= P_{12}, & Q_{12}Y_{12} &= P_{12}. \\ Z_{12}P_{12} &= Q_{12}, & Y_{12}P_{12} &= Q_{12}, \end{aligned}$$

and

$$P_{12}P_{23}P_{12} = Q_{12}Q_{23}Q_{12}.$$

The first sets of equations are now consistent and can be regarded simply as definitions of Y and Z . Using them in the second set we have

$$Y_{23}P_{23} = Q_{23} = Q_{12}^{-1}P_{12}P_{23}P_{12}Q_{12}^{-1} = Y_{12}P_{23}Y_{12}^{-1}.$$

We now write $P_{jk} = \hat{P}_{jk}\Phi_{jk}$, where Φ_{jk} is the flip which is the undeformed value of P . Then

$$Y_{23}\hat{P}_{23}\Phi_{23}Y_{12} = Y_{12}\hat{P}_{23}\Phi_{23},$$

and since $\Phi_{23}Y_{12} = Y_{13}\Phi_{23}$, this gives

$$Y_{23}\hat{P}_{23}Y_{13} = Y_{12}\hat{P}_{23}$$

Similarly we obtain

$$Y_{12}\hat{P}_{12}Y_{13} = Y_{23}\hat{P}_{12},$$

and from these we deduce that

$$(\hat{P}_{23}Y_{13}^{-1}\hat{P}_{23}^{-1})(\hat{P}_{12}Y_{13}^{-1}\hat{P}_{12}^{-1}) = (Y_{23}^{-1}Y_{12})(Y_{12}^{-1}Y_{23}) = 1,$$

and thence that

$$Y_{13}\hat{P}_{23}^{-1}\hat{P}_{12}Y_{13} = \hat{P}_{23}^{-1}\hat{P}_{12}.$$

Setting $V = \hat{P}_{23}^{-1}\hat{P}_{12}$, this can be rewritten as $Y_{13}VY_{13} = V$, which leads to

$$Y_{13}V^2 = Y_{13}VY_{13}VY_{13} = V^2Y_{13},$$

showing that Y_{13} and V^2 commute. Our standard square root argument shows that Y_{13} commutes with V , which in turn gives the identity $Y_{13}^2 = 1$, and so $Y_{13} = 1$, and $Q = P$. In the case in hand this shows that R commutes with Λ and so with Θ . ■

References

1. V. Chari & A. Pressley *A guide to Quantum Groups*, Cambridge University Press, 1994.
2. A. Connes, *Non-commutative geometry*, Academic Press, 1994.
3. A. Connes, M.R. Douglas and A.S. Schwarz, *Non-commutative geometry and matrix theory*, J. High Energy Phys. **2**, Paper 3 (1998), hep-th/9711162.
4. A. Kapustin, A. Kuznetsov, & D. Orlov, *Non-commutative instantons and twistor transform*, Comm. math. Phys. **221** (2001) 385-432, hep-th/0002193.
5. Yu.I. Manin, *Topics in non-commutative geometry*, Princeton UP, Princeton, 1991.
6. N. Nekrasov and A.S. Schwarz, *Instantons on non-commutative $\mathbf{R}^4$ and $(2,0)$ superconformal six-dimensional theory*, Comm. math. Phys. **198** (1998) 689-703, hep-th/9802068.
7. R. Penrose and W. Rindler, *Spinors in space-time, vol 2*, Cambridge University Press, 1986.
8. N. Seiberg and E. Witten, *String theory and non-commutative geometry*, J. High Energy Phys. **3**, Paper 32 (1999), hep-th/9908142.